

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
24 January 2002 (24.01.2002)

PCT

(10) International Publication Number
WO 02/06241 A1

(51) International Patent Classification: C07D 223/16,
A61K 31/55, A61P 11/08, 13/10, 1/12

Bramhall, Stockport, Cheshire SK7 2AE (GB). DAVIES,
Robin, Havard [GB/GB]; 12 Belgrave Court, 25 Cow-
bridge Road East, Cardiff CF1 9BJ (GB).

(21) International Application Number: PCT/GB01/02594

(22) International Filing Date: 14 June 2001 (14.06.2001)

(74) Agents: NASH, David, Allan et al.; Haseltine Lake &
Co., Imperial House, 15-19 Kingsway, London WC2B
6UD (GB).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0014561.5 14 June 2000 (14.06.2000) GB
0100600.6 9 January 2001 (09.01.2001) GB

(81) Designated States (national): AH, AG, AI, AM, AT, AU
(utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ, CZ (utility model), DE, DE
(utility model), DK, DK (utility model), DM, DZ, EC, EE,
EE (utility model), ES, FI, FI (utility model), GB, GI, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ,
LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,
MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI,
SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA, UG,
US, UZ, VN, YU, ZA, ZW.

(71) Applicant (for all designated States except US): MUSCA-
GEN LIMITED [GB/GB]; Morgan-Cole Solicitors,
Bradley Court, Park Place, Cardiff CF10 3DP (GB).



(72) Inventors; and

(84) Designated States (regional): ARIPO patent (GI, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,


(75) Inventors/Applicants (for US only): EVANS, Paul
[GB/GB]; Flat 10, 22 Gambier Terrace, Liverpool L1 7BL
(GB). THOMAS, Eric, James [GB/GB]; 6 Elmsway,

[Continued on next page]


(54) Title: 1, 2, 3, 5 -TETRAHYDROBENZO[C]AZEPIN-4-ONE DERIVATIVES HAVING MUSCARINIC ANTAGONIST AC-
TIVITY



(I)



(II)



(III)

(57) Abstract: There is disclosed a compound having the formula or a pharmaceutically acceptable salt thereof, wherein: R_{1a}, R_{1b} and R_{1c} are independently fluorine or hydrogen; R₂ is C₁ to C₁₂ alkyl being straight or branched chain, saturated or unsaturated, mono-substituted or unsubstituted, said substituents being selected from piperidine, pyrrolidine, morpholine, thiomorpholine and cycloalkyl of 3 to 7 carbon atoms; a cycloalkyl of 3 to 9 carbon atoms; a cycloalkyl of 3 to 9 carbon atoms having a C₁ to C₆ alkyl substituent; a polycycloalkyl of 2 to 3 rings having 7 to 12 carbons; and phenyl or phenyl substituted with halogen, hydroxy, C₁ to C₆ alkoxy, C₁ to C₆ alkyl, nitro, methylene dioxo or trifluoromethyl; and R₃ is a moiety selected from: (I), (II) or a pyrrolidine-3-yl moiety of the formula (III). The compounds are disclosed for use as muscarinic antagonists with M₃ selectivity.

WO 02/06241 A1



IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CH, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

with international search report

1,2,3,5-TETRAHYDROBENZO'CIAZEPIN-4-ONE DERIVATIVES HAVING MUSCARINIC ANTAGONIST ACTIVITY


THERAPEUTIC COMPOUNDS

This invention relates to muscarinic antagonists with M₃ selectivity.

Muscarinic M₃ receptors are located predominantly on smooth muscle and salivary glands, and agents selective for this sub-class of receptors may have therapeutic utility in the treatment of incontinence, disorders of gastro-intestinal motility and as bronchodilators in respiratory disease.


EP-A-0486734 discloses 1-substituted-1-hydroxy-1-aryl-3-(4-substituted-1-piperiziny)-2-propanones having antimuscarinic activity.

According to the present invention, there is provided a compound having the formula:




-2-

R_3 is a moiety selected from:




I



II

or a pyrrolidin-3-yl moiety of the formula



III

where R_6 is hydroxy or hydrogen;

where one of R_4 and R_5 is hydrogen or lower C1-3 alkyl and the other is selected from:

(a) hydrogen,
 (b) phenyl,
 (c) phenyl singly or multiply substituted with
 20 halogen, hydroxy, C_1 to C_6 alkoxy, C_1 to C_6 alkyl, nitro, methylene dioxy or trifluoromethyl; and

(d) C_1 to C_6 alkyl which may be branched chain or straight, saturated, unsaturated, or cyclic and may be optionally substituted with hydroxy, thienyl, pyrrolyl,
 30 pyridyl, furanyl, lower alkoxy or acetoxyalkyl wherein the alkyl group has 1 to 3 carbons, phenyl, phenyl singly or multiply substituted (preferably singly or doubly) with halogen, hydroxy, C_1 to C_6 alkoxy, C_1 to C_6 alkyl, nitro, methylene dioxy or trifluoromethyl.

35 In an embodiment of the invention, R_2 is not a

phenyl or substituted phenyl, R_3 has the structural formula II or III, and one of R_4 and R_5 is hydrogen whilst the other is selected from substituents (a), (b), (c) or (d).

5 Radical $R_{1a,b,c}$

In embodiments of the invention, R_{1a} , R_{1b} and R_{1c} are each fluorine or each hydrogen. In other embodiments, R_{1a} is hydrogen and either one of R_{1b} and R_{1c} is fluorine and the other is hydrogen or both R_{1b} and R_{1c} are
10 fluorine.

Radical R_2

When R_2 is substituted C_1 - C_{12} alkyl, the substituent on the alkyl may additionally be selected from tetrahydrofuran, thiophen and furan. Further, when R_2
15 is C_1 - C_{12} alkyl, it is preferred that the alkyl is saturated.

In preferred embodiments of the invention, R_2 may be cycloalkyl of 3 to 6 carbon atoms, for example cyclohexyl or cyclobutyl, preferably cyclobutyl. In
20 other preferred embodiments of the invention, R_2 may be phenyl.

Radicals R_4 and R_5

In addition to the above definition, under alternative (d) for R_4/R_5 , the or each alkyl substituent
25 on the phenyl radical may be a C_1 - C_{10} alkyl, preferably a C_5 - C_8 alkyl, and the or each alkoxy substituent on the phenyl radical may be C_1 - C_{10} alkoxy.

Further, in addition to the above definition under alternative (d) for R_4/R_5 , the methylene dioxy
30 substituent may itself be mono or di- substituted by an alkyl having 1 to 10 carbons, preferably dialkyl-substituted where each alkyl has from 1 to 5 carbons.

It is preferred that R_4 is hydrogen and R_5 is selected from amongst the groups (a)-(d) above.

35 In one embodiment, one of R_4 and R_5 is hydrogen (or

-4-

5 methyl in the case of R_5) and the other is selected from hydrogen, C_1 to C_6 alkyl which may be branched chain or straight, saturated, unsaturated, or cyclic and may be optionally substituted with hydroxy, thienyl, pyrrolyl, pyridyl, furanyl, phenyl, phenyl singly or multiply substituted (preferably singly or doubly) with halogen, hydroxy, C_1 to C_6 alkoxy, C_1 to C_6 alkyl or nitro. More preferably, one of R_4 and R_5 is hydrogen and the other is C_1 to C_6 alkyl, benzyl, substituted benzyl or
10 cinnamyl; such as benzyl or 4-substituted benzyl; for example benzyl, 4-chlorobenzyl or 4-methylbenzyl.


In another embodiment, it is preferred that R_4 is hydrogen and R_5 is C_1 to C_6 alkyl substituted by phenyl or phenyl which is singly or multiply substituted
15 (preferably singly or doubly) with halogen, hydroxy, C_1 to C_{10} alkoxy, C_1 to C_{10} alkyl or nitro. More preferably, R_5 is benzyl, substituted benzyl or cinnamyl. Most preferably, R_5 is substituted benzyl in which the substituent(s) on the benzyl are
20 independently halo, C_1 to C_{10} alkoxy or C_1 to C_{10} alkyl. For example, the benzyl may be substituted by one or two alkyls where the total number of carbon atoms in the alkyl substituent(s) is from 6 to 10. In another example, the benzyl may be substituted by an alkyl
25 radical having from 5-9 carbon atoms and a halo, preferably chloro. Where the benzyl is mono-substituted, this is preferably in the 3- or 4-position. Where the benzyl is di-substituted, this is preferably in the 3- and 4- positions.

30 Radical R_6

It is preferred that R_6 is hydrogen.

Preferred Embodiments of the Invention

In a first preferred embodiment of the invention, R_{1a} , R_{1b} and R_{1c} are independently hydrogen or fluorine,
35 R_2 is cycloalkyl of 3 to 6 carbon atoms or phenyl, R_3 is




where R_4 is hydrogen and R_5 is selected from C_1 to C_6 alkyl, benzyl, substituted benzyl or cinnamyl, and R_6 is hydrogen or hydroxy. Preferably R_6 is hydrogen.

In this first embodiment, R_2 is preferably cyclohexyl, cyclobutyl or phenyl, more preferably cyclobutyl, and R_5 is preferably C_1 to C_6 alkyl, benzyl, substituted benzyl or cinnamyl, such as methyl, benzyl or 4-substituted benzyl, for example benzyl, 4-chlorobenzyl or 4-methylbenzyl.

Alternatively in this first embodiment (and presently preferred), R_5 is substituted benzyl in which the substituent(s) on the benzyl are independently halo, C_1 to C_{10} alkoxy or C_1 to C_{10} alkyl. For example, the benzyl may be substituted by one or two alkyls where the total number of carbon atoms in the alkyl substituent(s) is from 6 to 10. In another example, the benzyl may be substituted by an alkyl radical having from 5-9 carbon atoms and a halo, preferably chloro. Where the benzyl is mono-substituted, this is preferably in the 3- or 4- position. Where the benzyl is di-substituted, this is preferably in the 3- and 4-positions.

In a second embodiment (which is presently less preferred than the first embodiment) R_{1a} , R_{1b} and R_{1c} are independently hydrogen or fluorine, R_2 is cycloalkyl of 3 to 6 carbon atoms or phenyl, R_3 is

- 6 -




5

where R_5 is hydrogen or methyl and R_4 is C_1 to C_6 alkyl, benzyl, substituted benzyl or cinnamyl, and R_6 is hydroxy or hydrogen, preferably hydrogen. In this second embodiment, R_2 is preferably cyclohexyl, cyclobutyl or phenyl, more preferably cyclobutyl, and R_4 is preferably C_1 to C_6 alkyl, benzyl, substituted benzyl or cinnamyl, such as methyl, benzyl or 4-substituted benzyl, for example benzyl, 4-chlorobenzyl or 4-methylbenzyl.

15

In a third embodiment R_{1a} , R_{1b} and R_{1c} are independently hydrogen or fluorine, R_2 is cycloalkyl of 3 to 6 carbon atoms or phenyl, R_3 is

20



25

where R_4 is hydrogen and R_5 is selected from C_1 to C_6 alkyl, benzyl, substituted benzyl or cinnamyl, and R_6 is hydroxy or hydrogen.

30

In this third embodiment, R_2 is preferably cyclohexyl, cyclobutyl or phenyl, more preferably cyclobutyl, and R_5 is preferably C_1 to C_6 alkyl, benzyl, substituted benzyl or cinnamyl, such as methyl, benzyl or 4-substituted benzyl, for example benzyl, 4-chlorobenzyl or 4-methylbenzyl.


35

Alternatively in this third embodiment (and presently preferred), R_5 is substituted benzyl in which the substituent(s) on the benzyl are independently

-7-

halo, C₁ to C₁₀ alkoxy or C₁ to C₁₀ alkyl. For example, the benzyl may be substituted by one or two alkyls where the total number of carbon atoms in the alkyl substituent(s) is from 6 to 10. In another example, the benzyl may be substituted by an alkyl radical having from 5-9 carbon atoms and a halo, preferably chloro. Where the benzyl is mono-substituted, this is preferably in the 3- or 4- position. Where the benzyl is di-substituted, this is preferably in the 3- and 4- positions.

In a fourth embodiment (which is presently less preferred than the third embodiment) R_{1a}, R_{1b} and R_{1c} are independently hydrogen or fluorine, R₂ is cycloalkyl of 3 to 6 carbon atoms or phenyl, R₃ is a pyrrolidin-3-yl moiety having the following structure:




where R₅ is hydrogen or methyl and R₄ is C₁ to C₆ alkyl, benzyl, substituted benzyl or cinnamyl, and R₆ is hydroxy. In this fourth embodiment, R₂ is preferably cyclohexyl, cyclobutyl or phenyl, more preferably cyclobutyl, and R₄ is preferably C₁ to C₆ alkyl, benzyl, substituted benzyl or cinnamyl, such as methyl, benzyl or 4-substituted benzyl, for example benzyl, 4-chlorobenzyl or 4-methylbenzyl.

In a fifth embodiment R_{1a}, R_{1b} and R_{1c} are independently hydrogen or fluorine, R₂ is cycloalkyl of 3 to 6 carbon atoms or phenyl, R₃ is a moiety having the following structure:

35

- 8 -



5


where R_4 is hydrogen and R_5 is selected from C_1 to C_6 alkyl, benzyl, substituted benzyl or cinnamyl.

10 In this fifth embodiment, R_2 is preferably cyclohexyl, cyclobutyl or phenyl, more preferably cyclobutyl, and R_5 is preferably C_1 to C_6 alkyl, benzyl, substituted benzyl or cinnamyl, such as methyl, benzyl or 4-substituted benzyl, for example benzyl, 4-chlorobenzyl or 4-methylbenzyl.

15 Alternatively in this fifth embodiment (and presently preferred), R_5 is substituted benzyl in which the substituent(s) on the benzyl are independently halo, C_1 to C_{10} alkoxy or C_1 to C_{10} alkyl. For example, the benzyl may be substituted by one or two alkyls
 20 where the total number of carbon atoms in the alkyl substituent(s) is from 6 to 10. In another example, the benzyl may be substituted by an alkyl radical having from 5-9 carbon atoms and a halo, preferably chloro. Where the benzyl is mono-substituted, this is
 25 preferably in the 3- or 4- position. Where the benzyl is di-substituted, this is preferably in the 3- and 4-positions.

In a sixth embodiment (which is presently less preferred than the fifth embodiment) R_{1a} , R_{1b} and R_{1c} are
 30 independently hydrogen or fluorine, R_2 is cycloalkyl of 3 to 6 carbon atoms or phenyl, R_3 is

- 9 -



5

where R₅ is hydrogen or methyl and R₄ is C₁ to C₆ alkyl, benzyl, substituted benzyl or cinnamyl. In this sixth embodiment, R₂ is preferably cyclohexyl, cyclobutyl or phenyl, more preferably cyclobutyl, and R₄ is preferably C₁ to C₆ alkyl, benzyl, substituted benzyl or cinnamyl, such as methyl, benzyl or 4-substituted benzyl, for example benzyl, 4-chlorobenzyl or 4-methylbenzyl.

In each of the first to sixth embodiments described above, it is preferred that R_{1a} is hydrogen and either one of R_{1b} and R_{1c} is fluorine and the other is hydrogen or both R_{1b} and R_{1c} are fluorine.

As used herein, unless otherwise specified, lower alkyl and lower alkoxy refer to groups having 1 to 6 carbons. The invention also relates to the pharmaceutically acceptable salts of the foregoing compounds and to pharmaceutical compositions containing effective amounts of such compounds; the compounds and compositions may be used for the manufacture of a medicament for the treatment of bladder disorders.

The compounds of the invention may be used in the neutral form. Alternatively, the compounds may be used in the form of pharmaceutically acceptable salts. Salts of the compounds of the invention include the acid salts such as the hydrochloride, sulfate, phosphate, nitrate, methanesulfonate and tartrate salts. Other pharmaceutically acceptable salts are also included in the invention, as are the various possible hydrates of each of the compounds. As will be understood by those skilled in the art, compounds of


-10-

this invention may be present as d or l optical isomers as well as racemic mixtures thereof. Further, some of the compounds in which R_1 is a substituted cycloalkyl or a polycycloalkyl may be present as diastereoisomers which may be resolved into optical isomers.

Resolutions of optical isomers may be accomplished by fractional crystallization of their salts with optically active acids such as, for example, tartaric, camphor-10-sulfonic, O,O-dibenzoyltartaric, O,O-di(p-toluoyl) tartaric, menthyloxyacetic, camphoric, or 2-pyrrolidone-5-carboxylic acids of N-acetyltryptophane from appropriate solvents. They may also be prepared by stereoselective synthesis or by chromatographic techniques using chiral substrates or derivatives.

Unless otherwise specified in the claims, it is intended to include all isomers, whether separated or mixtures thereof.

Preferred isomers have the following stereochemistry:



The protonated form of the respective R_3 side chains is shown.

The compounds of the invention may be administered in a variety of pharmaceutical preparations well known to those skilled in the pharmaceutical art. For

-11-

parenteral administration, the compounds may be prepared in aqueous injection solutions which may contain antioxidants, buffers, bacteriostats, and other additives commonly employed in such solutions.

5 Extemporaneous injection solutions may be prepared from sterile pills, granules or tablets which may contain diluents, dispersing and surface active agents, binders, and lubricants as well as the compound of the invention.


10 In the case of oral administration, fine powders or granules of the compound of the invention may be formulated with diluents and dispersing and surface active agents, and may be prepared in water, a syrup, capsules, cachets, a non-aqueous suspension or an
15 emulsion. In dry forms, optional binders and lubricants may be present. The compositions may also include flavorants, preservatives, suspending, thickening and emulsifying agents and other pharmaceutically acceptable additives. Granules or
20 tablets for oral administration may be coated. In general, the compositions of the invention include the compounds of the invention in a pharmaceutically effective amount in a pharmaceutically acceptable carrier.

25 The compounds are useful as antimuscarinic agents selective for the muscarinic M3 receptor; more particularly, they are useful as bronchodilators, as antispasmodics, antisecretory agents, have antiulcer activity and are useful in the treatment of patients
30 suffering from neurogenic bladder disorders. The compounds are administered in pharmaceutically effective amounts. Daily dosages will generally be at a rate of 5 to 100 mg/day, more specifically 10 to 40 mg/day. Because of their duration of action the
35 compounds may be administered less frequently than

certain prior art antimuscarinic agents, particularly those used in the treatment of neurogenic bladder disorder.


The compounds of the invention may be tested to determine their muscarinic activity in accordance with the procedure set forth in EP-A-0486734. The compounds may also be tested for their M_1 , M_2 and M_3 receptor activity using the assays set forth after the examples below.

According to a process aspect of the present invention, the compounds of the invention process may be synthesised by a process which includes the step of subjecting a compound of the formula (X):



in which R1a, R1b, R1c and R2 are as defined above and R3 is as defined above suitably protected, to oxidation conditions sufficient to oxidise the alcohol group at the 4-position of the benzo[c]azepine core to a ketone group.

For example, the R_3 groups may be protected as follows




-13-

in which Y is hydrogen or a hydroxy protecting group such as acetyl, and X is an amine protecting group such as a trifluoroacetamide or a nosyl group. In Formula III the nitrogen group only requires protection where R₅ in the final molecule is hydrogen.


The oxidation step to oxidise the alcohol group at the 4-position of the benzo[c]azepine core to a ketone group is preferably a Swern oxidation step (K. Takahashi, M. Ogata, J. Org. Chem, 1987, 52, 1877).

In this process aspect of the invention, the compound X may be made by a process in which a compound of the formula XI




XI


is subjected to a reductive amination with an aldehyde/ketone corresponding to R₃, suitably protected. For example, one of the following protected aldehyde or ketone may be employed:



Ia



IIa



IIIa

where X and Y are as defined above.

This reductive amination may be accomplished

-14-

following the procedure of Borch et al. (R.F. Borch, M.D. Bernstein, H.D. Hurst, J. Am. Chem. Soc., 1971, 93, 2897) using the reagent NaBH_3CN at an optimum pH of about 6.

5 Details of the routes to the compound XI and reagents Ia, IIa and IIIa are discussed in detail below.


10 The following section concerns the synthesis of compounds in accordance with the invention in which each of R_{1a} , R_{1b} and R_{1c} is hydrogen, and uses as starting material the commercially available compound phthalide (isobenzofuran).

15

20

-15-

Synthesis of 2-Azepine Core Unit:



Ketone 5 is accessed in two ways via the Weinreb amide 2 or via an addition-oxidation protocol. 6 is prepared by treatment of 5 with the Petasis reagent (Cp_2TiMe_2).


- 5 Deprotection, oxidation followed by a two step reductive amination gives the dialkenyl amine 9. Nitrogen derivatisation with 2-nitrophenyl sulfonyl chloride gives 10 which is converted to the dihydroazepine 11 using tricyclohexylphosphine-[1,3-bis(2,4,6-trimethylphenyl)-4,5-dihydroimidazol-2-ylidene] [benzylidene]ruthenium dichloride. This ring closing step follows the methodology developed by Grubbs using catalysts based on ruthenium (P. Schwab,
- 10

-16-

R.H. Grubbs, J.W. Ziller, J. Am. Chem. Soc., 1996, 118, 100; S.T. Nguyen, R.H. Grubbs, J.W. Ziller, J. Am. Chem. Soc., 1993, 115, 9858; E.L.Dias, S.T. Nguyen, R.H. Grubbs, J. Am. Chem. Soc., 1998, 63, 824). Other metal catalysts for the construction of cyclic amines are known in the art. Dihydroxylation using OsO_4 gives the diol 12. The sulfonyl group in 12 is removed using thiophenol giving the amino diol 14. In the first scheme the reductive coupling of butyraldehyde to give 15 is shown, although outside the scope of the invention. The coupling of different side chains is described in more detail below. Swern oxidation generates the β -amino ketone 16.

The aldehydes and ketones containing the second amino group were synthesised as follows:

C-3 Side Chain 20:




Two step reductive amination between benzaldehyde 17a and but-4-enylamine gives the amine 18. Protection of 18 as the trifluoroacetamide gives the alkene 19, treatment with ozone gives the aldehyde 20 on reductive work-up.

The hydroxy C3 side chain may be synthesised as indicated below with an asymmetric hydroxylation providing enantiomerically enriched material. The key steps are the reductive amination of the aryl aldehyde with allylamine 201 to give the secondary amine 202. Protection of this either as its *o*-nitrobenzenesulfonyl or trifluoroacetyl derivative 203 followed by asymmetric hydroxylation to provide the diol 204. Regioselective *O*-silylation and then acetylation of the

-17-


secondary alcohol will provide the acetate 205 which on desilylation and oxidation will give the required aldehyde 206. Reductive coupling of the aldehyde then proceeds as described above.

5



10

C-2 Side Chains 25a and 25b



15

Following a similar sequence described above reductive amination of benzaldehyde 17a and 2-naphthylaldehyde 17b with the acetal 21 gives the amines 22a and 22b. Protection of the amino group with trifluoroacetic anhydride gives the amides 23a and 22b. Acetal deprotection under acidic catalysis furnishes the aldehydes 24a and 24b.

20

Pyrrolidine Side Chains 31 and 33:


Reference is made to the following reaction scheme.

25

Trans-4-Hydroxyproline 25 is protected with the tert-butyloxy carbonyl group (Boc) 26 before acid activation as the Weinreb amide and hydroxyl protection as the tert-butyldimethylsilyl ether (TBS) giving the known amide 27. Organometallic addition to 27 with

-18-


phenyl magnesium bromide (PhMgBr) gives the ketone 28. Reduction of the amide 27 with diisobutylaluminium hydride gives the aldehyde 29. Carbonyl reduction of 28 and 29 generates the protected pyrrolidines 30 and 31. TBS removal followed by Swern oxidation generates the Boc protected ketones 31 and 33.



An alternative synthesis of the pyrrolidine side chains is as follows.

With reference to the following reaction scheme, the starting material hydroxyproline 101 is bis-protected and converted into the ester 102. Reduction - oxidation to the aldehyde 103 followed by addition of an aryl Grignard reagent will give the alcohol 104, as a mixture of diastereoisomers, which is reduced using Barton chemistry (e.g. conversion into the thionocarbonate followed by reduction using tributyltin hydride) to give the pyrrolidine derivative 105. At this stage the t-Boc group will be replaced by a trifluoroacetate and desilylation and oxidation will give the ketone 106. Reductive coupling of the ketone then proceeds as described above.


-19-



Side Chain Coupling to Azepinyl Nucleus 14:


15 The side chains whose synthesis is described above are then coupled to the functionalised azepinyl nucleus 14 under one-pot reductive amination conditions. This furnishes the diols 34a-c and 36a,b. Swern oxidation of the secondary hydroxyl group and deprotection of the trifluoroacetamide 34a-c, or tert-butyloxycarbonyl

20 35a,b gives the azepines 35a-c and 37a,b.



The 5-fluoro analogues may be synthesised as follows:

Synthesis of 5-Fluoro Analogue:



Phthalide **1** is regio-selectively nitrated to give **38**.

Reduction of the nitro group gives the aniline **39**,

5 which is converted into the known 5-fluorophthalide **40**.

Diisobutylaluminium hydride reduction of the lactone in **40** gives the lactol **41**. Reductive amination of **41**

with allylamine gives the amine **42**, which is


chemo-selectively converted to its

-21-

2-nitrophenylsulfonyl derivative 43. Manganese dioxide oxidation of 43 gives the aldehyde 44 which reacts with cyclobutylmagnesium bromide to give the benzylic alcohol 45. Oxidation giving the ketone 46 followed by methylenation with Petasis's reagent gives the dialkenyl cyclisation precursor 47. Ring closing olefin metathesis with the imidazolyl based ruthenium benzylidene catalyst gives the dihydroazepine 48. Dihydroxylation with OsO₄ gives the diol 49, which is converted into the amine 50. 50 is then coupled with the acyclic side chains 20 and 24a,b under the one-pot reductive amination conditions to give the diols 51a-c. Oxidation, followed by trifluoroacetamide deprotection gives the 4-fluoro bis-amines 52a-c.


An alternative (less preferred) synthetic pathway to that described above is as follows. These pathways may be generalized by the skilled person where necessary.

For compounds where R₃ is as follows:



the following synthetic pathway may be employed.

-22-



Step a Reaction of the cyclobutyl lithium (or other organometallics) with the known lactone is conducted at -78°C , with slow inverse addition of the reagent.

-23-

Step b Reaction of the ketoalcohol with bromoacetyl bromide and pyridine (mole ratio 1:1:1), gives an unstable bromoester; which is utilised immediately.

5 **Steps c** The bromoester is dissolved in acetonitrile at room temperature and treated with triphenylphosphine. After 3 days stirring at room temperature triethylamine is added and after a further week the ϵ -lactone is isolated.

10 **Steps d** The lactone may be reduced in one step, but the better yields are achieved by use of a two step protocol via the lactol.

15 **Steps e** The bis(tosylate) is prepared at -20°C warming to room temperature to minimise formation of a cyclic ether. Formation of the seven membered ring is performed under high dilution conditions (0.1 mmolar) in DMSO to minimise dimer formation. The methodology is illustrated by the oxazolidinone protected amino alcohol, but other protecting groups may be employed such as carbonenzyloxy.


20 **Steps f** The diol moiety is installed by use of Sharpless asymmetric dihydroxylation methodology using the AD-mix α or a comparable reagent.

Steps g Oxidation of the diol is performed under mild conditions using the Dess-Martin periodinane reagent.

25 **Steps h** The oxazolidinone ring is cleaved directly under acidic or basic conditions. In the case where $R = \text{H}$, the three step method indicated in the scheme is the preferred method.

For compounds in which R_3 is as follows:


30



35 the same pathway as given above in the immediately

-24-

preceding section may be used except that a different side chain is used in the N-alkylation step (step (e)).




The invention will now be illustrated by the following examples.

In the examples, flash column chromatography was performed using Merck silica gel (60H; 40-60 μ , 230-240 mesh). Thin layer chromatography (TLC) was carried out using glass backed plates coated with Merck HF254/366 silica gel. The spots were visualised using ultraviolet radiation, treatment with basic permanganate solution, or acidic ethanolic anisaldehyde solution. Petroleum ether (Pet) was redistilled before use and refers to the fraction boiling between 40 and 60°C. Tetrahydrofuran was dried over sodium-benzophenone and was distilled prior to use. Dichloromethane was dried over CaH₂ and was distilled before use.

Mass spectra, either electron impact (EI), or chemical ionisation using ammonia (CI), were recorded by Val Boote using a Fisons VG Trio 200 spectrometer. High resolution mass spectra were recorded by Peter Kobryn on a Kratos Concept IS spectrometer. Microanalyses were performed using a Carlo-Erba combustion analyser for C, H and N. Infra-red spectra were recorded on a Genesis FTIR spectrometer on NaBr plates, either neat, or as evaporated films. Proton, proton-decoupled carbon and fluorine nuclear magnetic resonance spectra were recorded on either a Varian (400 MHz), Varian INOVA 300 (300 MHz), or a Varian Gemini 200 (200 MHz) spectrometer. Where applicable proton assignment was facilitated using correlation spectroscopy (COSY). Residual non-deuterated solvent was used as an internal standard and the chemical shifts are quoted in ppm down field from tetramethylsilane. Signal splitting patterns are

described as singlets (s), doublets (d), doublet of doublets (dd), doublet of double doublets (ddd), triplets (t), doublet of triplets (dt), quartets (q), or multiplets (m). The coupling constants (J) are given in Hertz (Hz).

Example 1




N-Benzyl-N-(3-butenyl)amine 1:

- 10 According to literature⁷⁹ at 0°C AlCl₃ (18.0 g, 0.135 mol, 1 eq.) in dry Et₂O (200 cm³) was treated initially with LAH (5.12 g, 0.135 mol, 1 eq.) and then after 0.5h allyl cyanide (9.3 g, 0.135 mol, 1 eq.) was added dropwise. Stirring was maintained for 2 h at 0°C
- 15 before H₂O (20 cm³) was added followed by 4 M NaOH (20 cm³) and H₂O (60 cm³). The solid residue was filtered, washing with Et₂O (2 x 50 cm³). The volatile amine was stripped *in vacuo* with care and added directly to a solution of benzaldehyde (14 cm³, 0.137 mol, 1.01 eq.)
- 20 in DCM (200 cm³) with MgSO₄ (20 g). Stirring was continued at room temperature for 24 h. Filtration followed by solvent removal gave the imine, which was reduced directly. The imine in MeOH (100 cm³) was treated portionwise with NaBH₄ (5.1 g, 0.134 mol, 1
- 25 eq.) and stirring was continued for 2 h. The reaction mixture was concentrated *in vacuo* and Et₂O (100 cm³) and H₂O (100 cm³) were added. The resultant aqueous phase was further extracted with Et₂O (2 x 100 cm³) and the combined organic phases were dried over MgSO₄.
- 30 Filtration and solvent removal under reduced pressure


gave 1 (5.9 g, 44%) as a yellow liquid. m/z (CI) 162 (MNH_4^+ , 100%); δ_H (300 MHz, $CDCl_3$) 2.25 (2H, q, J 7.0 Hz, CH_2), 2.76 (2H, t, J 7.0 Hz, CH_2), 3.82 (2H, s, CH_2), 5.02-5.19 (2H, m, CH_2), 5.76-5.94 (1H, m, CH), 7.23-7.44 (5H, m, ArH).

Example 2




N-Benzyl-N-(3-butenyl)-2,2,2-trifluoroacetamide 2:

At 0°C a solution of amine 1 (5.9 g, 36.65 mmol, 1 eq.) and TEA (25.0 cm³, 179.37 mmol, 5 eq.) in DCM (100 cm³) was treated with a solution of (CF₃CO)₂O (7.8 cm³, 55.22 mmol, 1.5 eq.) added via a dropping funnel. The mixture was stirred for 3 h at 0°C to room temperature. HO (100 cm) was added and the resultant aqueous phase was further extracted with DCM (2 x 100 cm³). The combined organic extracts were dried over MgSO₄. Filtration, solvent removal under reduced pressure and purification by flash column chromatography (Pet:Et₂O; 9:1) gave 2 (6.42 g, 68%) as a yellow liquid. R_f = 0.3 (Pet:Et₂O; 9:1); m/z (CI) 275 (MNH_4^+ , 100%), 258 (MH^+ , 40%); found (EI) 257.1022, C₁₃H₁₄NOF₃ requires 257.1027 (-1.9 ppm); δ_H (300 MHz, $CDCl_3$) 2.23-2.42 (2H, m, CH_2), 3.34-3.45 (2H, m, CH_2), 4.62 (2H, s, CH_2), 4.71 (2H, s, CH_2), 5.05-5.18 (2H, m, CH_2), 5.63-5.81 (1H, m, CH), 7.11-7.22 (5H, m, ArH); ¹H-NMR spectrum complicated due to restricted rotation.

Example 3

N-Benzyl-N-(3-oxopropyl)-2,2,2-trifluoroacetamide 3:


- At -78°C a solution of 2 (2.01 g, 7.82 mmol, 1 eq.) in
 5 DCM (20 cm³) was treated with a steady stream of ozone
 gas for 0.5 h. At this point TLC analysis indicated
 consumption of 2. The excess ozone was purged under a
 flow of oxygen and DMS (3 cm, 40.86 mmol, 5.2 eq.) was
 added. The reaction mixture was warmed to room
 10 temperature and stirred for 15 h. Solvent removal in
 vacuo and flash column chromatography (Pet:EtOAc; 4:1,
 1% TEA) gave 3 (1.72 g, 85%) as a clear liquid. R_f =
 0.2 (Pet:EtOAc; 4:1); ν_{\max} (neat/cm⁻¹) 3066, 3034,
 2944, 2836, 2733, 1724, 1690, 1453, 1377, 1204, 1147;
 15 m/z (CI) 277 (MNH₄⁺, 100%), 260 (MH⁺, 100%); found
 277.1167, C₁₂H₁₂NO₂F₃·NH₄ requires 277.1164 (+1.1 ppm);
 δ_H (300 MHz, CDCl₃) 2.75-2.84 (2H, m, CH₂), 3.61 (2H,
 t, J 6.5 Hz, CH₂), 3.76 (2H, t, J 7.0 Hz, CH₂), 4.66
 (2H, s, CH₂), 4.73 (2H, s, CH₂), 7.25-7.47 (5H, m,
 20 ArH), 9.74 (1H, s(br), CHO); ¹H-NMR spectrum
 complicated due to restricted rotation (60:40).

Example 4

- 25 N-Allyl-N-benzylamine 4:
 A mixture of benzaldehyde (9.8 cm³, 96.0 mmol, 1 eq.),
 allylamine (10.8 cm³, 143.9 mmol, 1.5 eq.) and MgSO₄
 (20 g) in DCM (100 cm³) were stirred together at room
 temperature for 15 h. Filtration and solvent removal

under reduced pressure gave the imine (ca. 96 mmol) which was dissolved in MeOH (100 cm³). At room temperature NaBH₄ (3.65 g, 96.1 mmol, 1 eq.) was added in portions. Stirring was continued for 2 h before
5 approximately half the solvent was removed under reduced pressure. Et₂O (100 cm³) and H₂O (100 cm³) were added and the mixture was basified with 1 M NaOH (ca. pH 12). The aqueous layer was further extracted with Et₂O (2 x 100 cm³) and the combined ethereal
10 extracts were dried over MgSO₄. Filtration and solvent removal in vacuo afforded the amine 4 (8.94 g, 64%) as a clear liquid. m/z (CI) 148 (MH⁺, 100%); δ _H (200 MHz, CDCl₃) 1.5 (1H, s(br), NH), 3.32 (2H, d, J 6.5 Hz, CH₂), 4.82 (2H, s, CH₂), 5.07-5.32 (2H, m, CH₂), 5.82-
15 6.08 (1H, m, CH), 7.19-7.45 (5H, m, ArH).

Example 5




N-Allyl-N-benzyl-2-nitrophenylsulfonamide 5:

20 At room temperature a mixture of 4 (950 mg, 6.46 mmol, 1 eq.), TEA (1.8 cm³, 12.91 mmol, 2 eq.) and DMAP (ca. 2 mg) in DCM (20 cm³) was treated with NsCl (1.43 g, 6.45 mmol, 1 eq.). Stirring was continued for 3 h before H₂O (50 cm³) and Et₂O (50 cm³) were added. The
25 resultant aqueous phase was extracted further with Et₂O (2 x 50 cm³) and the total organic extracts were dried over MgSO₄. Filtration followed by solvent removal in vacuo afforded the crude sulfonamide which was purified by flash column chromatography (Pet:EtOAc; 5:1) to give

5 (1.63 g, 76%) as a viscous clear oil. $R_f = 0.25$
(Pet:EtOAc; 5:1); m/z (CI) 350 (MNH_4^+ , 100%); found
350.1176, $C_{16}H_{16}N_2O_4S \cdot NH_4$ requires 350.1174 (+0.6 ppm);
 δ_H (300 MHz, $CDCl_3$) 3.77 (2H, d, J 6.5 Hz, CH_2), 4.45
5 (2H, s, CH_2), 4.95–5.08 (2H, m, CH_2), 5.50 (1H, ddd
app. qt, J 6.5, 10.0, 17.0 Hz, CH), 7.19–7.25 (5H, m,
ArH), 7.51–7.65 (3H, m, ArH), 7.93 (1H, d, J 7.5 Hz,
ArH); δ_C (75 MHz, $CDCl_3$) 49.2, 50.3, 119.6, 124.2,
127.8, 128.3, 128.6, 130.9, 131.7, 131.8, 133.5, 134.0,
10 135.3, 147.8.

Example 6




N-Benzyl-N-(2-oxoethyl)-2-nitrophenylsulfonamide 6:

15 A solution of 5 (1.11 g, 3.34 mmol, 1 eq.) in DCM (25
cm³) at -78°C was treated with a steady stream of ozone
gas until TLC analysis indicated no remaining starting
material (ca. 0.5 h). The excess ozone was purged
under a flow of oxygen before DMS. (4.0 cm³, 54.47 mmol,
20 16 eq.) was added and the mixture was allowed to warm
to room temperature and stirred for 15 h. Evaporation
of the solvent under reduced pressure followed by flash
column chromatography (Pet:EtOAc; 1:1) gave 6 (1.04 g,
92 %) as a colourless solid. For microanalysis 6 was
25 recrystallised from EtOAc and petroleum ether. $R_f =$
0.25 (streak) (Pet:EtOAc; 5:1); ν_{max} ($CDCl_3/cm^{-1}$) 3055,
2986, 2831, 1735, 1546, 1371, 1266, 1166; m/z (FAB) 690
(M_2Na^+ , 90%); δ_H (300 MHz, $CDCl_3$) 4.11 (2H, s, CH_2),
4.65 (2H, s, CH_2), 7.25–7.38 (5H, m, ArH), 7.65–7.74

(3H, m, ArH), 8.11 (1H, d, J 7.5 Hz, ArH), 9.39 (1H, s, CHO); δ_C (75 MHz, CDCl₃) 52.6, 55.5, 124.3, 128.6, 128.7, 128.9, 130.9, 131.9, 133.0, 133.8, 134.1, 147.0, 196.4; found C, 53.6; H, 4.5; N, 8.4; S, 9.6%;
5 C₁₅H₁₄N₂O₅S requires, C, 53.9; H, 4.2; N, 8.4; S, 9.6%.


Example 7



2-(tert-Butyldimethylsilyloxymethyl)-N-methoxy-N-
10 methylbenzamide 7:
At 0°C under argon a 2 M solution of AlMe₃ in hexane
(32 cm³, 64.5 mmol, 2 eq.) was added dropwise over ca.
0.25 h to a suspension of HCl·NH(OMe)Me (6.29 g, 64.5
mmol, 2 eq.) in DCM (60 cm³). During the addition of
15 0.5 eq. of AlMe₃ a vigorous gas evolution ensued. The
now clear mixture was stirred for 1 h at 0°C before a
solution of phthalide (isobenzofuran) (4.32 g, 32.2
mmol, 1 eq.) in DCM (20 cm³) was added. Stirring was
maintained for 7 h during which time room temperature
20 was reached. Saturated sodium potassium tartrate
solution (100 cm³) was cautiously added. The resultant
aqueous layer, obtained after separation was further
extracted with DCM (2 x 100 cm³). The combined organic
extracts were washed with sat. brine solution (100 cm³)
25 and dried over Na₂SO₄. Filtration followed by solvent
removal in vacuo gave the crude Weinreb amide which was
directly O-protected in order to minimise re-
lactonisation. Thus, at room temperature the crude
amide (ca. 32.2 mmol, 1 eq.) was dissolved in DCM (50
30 cm³) and treated with TBDMS-Cl (4.9 g, 32.2 mmol, 1

eq.) and imidazole (4.4 g, 64.5 mmol, 2 eq.). Stirring was continued for 15 h. Water (100 cm³) and DCM (50 cm³) were added and the resultant aqueous layer was further extracted with DCM (100 cm³). The combined organic extracts were dried over MgSO₄ before filtration and solvent removal in vacuo gave the crude product. Purification by flash column chromatography (Pet:EtOAc; 5:1 → Pet:EtOAc; 3:1) gave the Weinreb amide 7 (5.93 g, 60%) as a clear liquid. $R_f = 0.3$ (Pet:EtOAc; 4:1); ν_{\max} (neat/cm⁻¹) 3064, 2954, 2893, 2857, 1650, 1463, 1416, 1383, 1257, 1119, 1081; m/z (CI) 310 (MH⁺, 100%); found 310.1835, C₁₆H₂₇NO₃Si·H requires 310.1838 (+0.9 ppm); δ_H (300 MHz, CDCl₃) 0.00 (6H, s, CH₃), 0.84 (9H, s, CH₃), 3.19 (3H, s, CH₃), 3.44 (3H, s(br), CH₃), 4.69 (2H, s, CH₂), 7.16–7.22 (2H, m, ArH), 7.31 (1H, dt, J 2.5, 7.5 Hz, ArH), 7.45 (1H, d, J 7.5 Hz, ArH); δ_C (75 MHz, CDCl₃) -5.4, 18.4, 25.9, 33.6, 61.0, 62.5, 126.4, 126.8, 129.3, 132.7, 138.8, 169.7.

20


Example 8

(2-tert-Butyldimethylsilanyloxymethyl)phenyl]cyclobutylmethanone 8:

25 At -78°C under argon ^tBuLi 1.7 M in pentane (16 cm³, 27.31 mmol, 2 eq.) was added in a dropwise fashion to a solution of cyclobutyl bromide (1.3 cm³, 13.66 mmol, 1 eq.) in THF (15 cm³). The resultant yellow solution was stirred for 1 h at -78°C before adding via cannula

to a cooled (-78°C) solution of the Weinreb amide 7 (2.11 g, 6.83 mmol, 0.5 eq.) in THF (30 cm^3). Stirring was continued for 1 h. Saturated NH_4Cl solution (50 cm^3) was added and the mixture was warmed to room temperature. Extraction with ether ($3 \times 50\text{ cm}^3$) and drying of the combined organic extracts over MgSO_4 gave the crude cyclobutane after filtration and solvent removal under reduced pressure. Purification by flash column chromatography (Pet:EtOAc; 8:1 \rightarrow Pet:EtOAc; 5:1) afforded **8** (1.1 g, 59%) as a clear oil. $R_f = 0.55$ (Pet:EtOAc; 5:1).

Example 9




tert-Butyl-[2-(1-cyclobutylvinyl)benzyloxy]dimethylsilane **9**:

Under N_2 in foil covered apparatus a mixture of **8** (476 mg, 1.57 mmol, 1 eq.) and $\text{Cp}_2\text{TiMe}_2^{73}$ (700 mg, 3.35 mmol, 2.1 eq.) in THF (20 cm^3) were heated to reflux for 15 h. Petroleum ether (100 cm^3) was added and the reaction mixture was filtered through Celite®. The residue was washed with petroleum ether ($2 \times 50\text{ cm}^3$) before silica (ca. 5 g) was added and the solvent removed under reduced pressure. Purification by flash column chromatography (Pet:EtOAc; 19:1) gave **9** (434 mg, 92%) as a clear oil. $R_f = 0.25$ (Pet:EtOAc; 19:1); m/z (CI) 320 (MNH_4^+ , 5%), 303 (MH^+ , 10%), 171 (100%); found (EI) 302.2062 $\text{C}_{19}\text{H}_{30}\text{OSi}$ requires 302.2066 (-1.3 ppm);

δ_H (300 MHz, $CDCl_3$) -0.08 (6H, s, CH_3), 0.84 (9H, s, CH_3), 1.55-1.67 (1H, m, CH_2), 1.69-2.01 (5H, m, CH_2), 3.10 (1H, pent, J 8.0 Hz, CH), 4.58 (2H, s, CH_2), 4.79 (1H, d, J 1.5 Hz, CH_2), 5.06 (1H, d, J 1.5 Hz, CH_2),
5 6.94 (1H, d, J 7.5 Hz, ArH), 7.09 (1H, t, J 7.5 Hz, ArH), 7.17 (1H, t, J 7.5 Hz, ArH), 7.44 (1H, d, J 7.5 Hz, ArH); δ_C (75 MHz, $CDCl_3$) -5.3, 17.6, 18.4, 25.9, 28.0, 42.0, 62.6, 112.1, 126.2, 126.7, 127.9, 138.0, 140.2, 151.9.


10

Example 10

[2-(1-Cyclobutylvinyl)phenyl]methanol 10:


At room temperature a 1 M solution of TBAF (0.9 cm³,
15 0.90 mmol, 1 eq.) was added dropwise to a solution of 9 (269 mg, 0.89 mmol, 1 eq.) in THF (10 cm³) and stirring was continued for 2 h. Et₂O (15 cm³) and H₂O (25 cm³) were added and the resultant aqueous layer was extracted with Et₂O (2 x 25 cm³). The combined
20 ethereal extracts were dried over MgSO₄, filtered and the solvent removed under reduced pressure. Purification by flash column chromatography (Pet:EtOAc; 19:1 → Pet:EtOAc; 5:1) afforded the title compound 10 (139 mg, 83%) as a clear oil. R_f = 0.3 (Pet:EtOAc;
25 5:1); m/z (CI) 206 (MNH_4^+ , 15%), 189 (MH^+ , 10%), 171 (100%); found 189.1277, $C_{13}H_{16}O \cdot H$ requires 189.1279 (-1.1 ppm); δ_H (300 MHz, $CDCl_3$) 1.55-1.67 (1H, m, CH_2), 1.71-2.01 (5H, m, CH_2), 3.13 (1H, pent, J 8.0 Hz, CH),

4.54 (2H, s, CH₂), 4.83 (2H, d, *J* 1.25 Hz, CH₂), 5.11 (1H, dd, *J* 1.25 Hz, CH₂), 6.99 (1H, d, *J* 7.0 Hz, ArH), 7.11-7.22 (2H, m, ArH), 7.36 (1H, d, *J* 7.25 Hz, ArH); δ_C (75 MHz, CDCl₃) 17.6, 27.9, 42.1, 63.2, 112.4, 127.05, 127.1, 128.0, 128.5, 137.7, 141.7, 152.2.


Example 11

2-(1-Cyclobutylvinyl)benzaldehyde 11:

10 A solution of 10 (1.90 g, 10.11 mol, 1 eq.) in DCM (60 cm³) was treated with pre-dried MnO₂ (4.40 g, 50.61 mol, 5 eq.) at room temperature and stirring was continued for 2 days. The reaction mixture was then filtered through Celite® and the residue was washed with DCM (2 x 50 cm³). Concentration in vacuo and purification by flash column chromatography (Pet:EtOAc; 19:1) gave the aldehyde 11 (1.75 g, 93%) as a clear liquid. *R_f* = 0.3 (Pet:EtOAc; 19:1); ν_{\max} (neat, cm⁻¹) 3084, 2976, 2940, 2864, 2748, 1695, 1596, 1479, 1446, 1391; *m/z* (CI) 264 (MNH₄⁺, 15%), 187 (MH⁺, 60%), 169 (100%); found 187.1121, C₁₃H₁₄O·H requires 187.1122 (-0.5 ppm); δ_H (300 MHz, CDCl₃) 1.66-1.77 (1H, m, CH₂), 1.79-2.13 (5H, m, CH₂), 3.31 (1H, pent, *J* 8.0 Hz, CH), 4.96 (1H, d, *J* 1.5 Hz, CH₂), 5.37 (1H, d, *J* 1.5 Hz, CH₂), 7.25 (1H, d, *J* 7.5 Hz, ArH), 7.37 (1H, t, *J* 7.5 Hz, ArH), 7.52 (1H, t, *J* 7.5 Hz, ArH), 7.93 (1H, d, *J* 7.5 Hz, ArH), 10.18 (1H, s, CHO); δ_C (75 MHz, CDCl₃) 17.6, 27.7, 42.2, 115.5, 127.2, 127.3, 128.9, 133.2, 133.7, 146.2, 149.2, 192.1.

Example 12*Allyl-[2-(1-cyclobutylvinyl)benzyl]amine 12:*

5 A mixture of aldehyde 11 (540 mg, 2.90 mmol, 1 eq.) and
MgSO₄ (ca. 5 g) in DCM (30 cm³) were treated at room
temperature with allylamine (0.45 cm³, 6.00 mmol, 2
eq.). The reaction mixture was stirred for 24 h and
filtered. Solvent removal gave the crude imine. At
10 room temperature the imine (ca. 2.90 mmol, 1 eq.) was
dissolved in MeOH (20 cm³) and NaBH₄ (164 mg, 4.33
mmol, 1.5 eq.) was added portionwise. After stirring
for 2 h DCM (50 cm³) and H₂O (50 cm³) were added and
the mixture was basified with 2.5 M NaOH (pH 10). The
15 resultant aqueous phase was further extracted with DCM
(3 x 50 cm³) and the combined organics were dried over
MgSO₄. Filtration and solvent removal *in vacuo* gave
127 (600 mg, 91%) as a yellow oil which was used
without further purification. m/z (CI) 288 (MH⁺,
20 100%); δ_{H} (300 MHz, CDCl₃) 1.39-1.51 (1H, s(br), NH),
1.68-1.79 (1H, m, CH₂), 1.87-2.11 (5H, m, CH₂), 3.21
(1H, pent, *J* 8.0 Hz, CH), 3.25 (2H, dt, *J* 0.5, 6.5 Hz,
CH₂), 3.75 (2H, s, CH₂), 4.93 (1H, s, CH₂), 5.07-5.25
(2H, m, CH₂), 5.19 (1H, s, CH₂), 5.84-6.03 (1H, m, CH),
25 7.04-7.08 (1H, m, ArH), 7.16-7.29 (2H, m, ArH), 7.39-
7.43 (1H, m, ArH); δ_{C} (75 MHz, CDCl₃) 17.6, 28.0, 42.2,
50.6, 51.9, 53.3, 112.0, 115.6, 126.3, 126.8, 128.6,
128.7, 136.9, 137.0, 141.8, 152.7.


Example 13

N-Allyl-N-[2-(1-cyclobutylvinyl)benzyl]-2-nitrophenylsulfonamide 13:

- 5 A mixture of the amine 12 (200 mg, 0.881 mmol, 1 eq.), TEA (0.18 cm³, 1.291 mmol, 1.5 eq.), 2-NsCl (215 mg, 0.970 mmol, 1.1 eq.) and a catalytic amount of DMAP (ca. 2 mg) in DCM (10 cm³) were stirred at room temperature for 3 h. Et₂O (25 cm³) and H₂O (25 cm³)
- 10 were added and the resultant aqueous layer was further extracted with Et₂O (2 x 15 cm³). The combined organic extracts were dried over MgSO₄. Filtration, solvent removal under reduced pressure followed by flash column chromatography (Pet:EtOAc; 3:1) gave the title compound
- 15 13 (285 mg, 79%) as a clear viscous oil. *R*_F = 0.3 (Pet:EtOAc; 3:1); *m/z* (CI) 430 (MNH₄⁺, 15%), 413 (MH⁺, 5%), 383 (30%), 228 (50%), 171 (100%); found 413.1528, C₂₂H₂₄N₂O₄S·H requires 413.1535 (-1.7 ppm); δ _H (300 MHz, CDCl₃) 1.63-1.78 (1H, m, CH₂), 1.82-2.11 (5H, m, CH₂), 3.16 (1H, pent, *J* 8.25 Hz, CH), 3.94 (2H, d, *J* 6.25 Hz, CH₂), 4.61 (2H, s, CH₂), 4.88 (1H, s, CH₂), 5.01-5.12 (2H, m, CH₂), 5.22 (1H, t(br), *J* 1.5 Hz, CH₂), 5.61 (1H, tq, *J* 6.25 Hz, CH), 7.04-7.11 (1H, m, ArH), 7.18-7.26 (2H, m, ArH), 7.35-7.41 (1H, m, ArH),
- 20 7.63-7.76 (3H, m, ArH), 8.04 (1H, d, *J* 7.5 Hz, ArH); δ _C (75 MHz, CDCl₃) 17.6, 28.0, 42.0, 48.1, 49.6, 112.8,
- 25

119.0, 124.1, 126.8, 127.1, 127.2, 128.6, 131.0, 131.6,
132.0, 132.2, 133.4, 134.0, 141.8, 143.8, 151.6.

Example 14




5 *5-Cyclobutyl-2-(2-nitrophenylsulfonyl)-2,3-dihydro-1H-benzo[c]azepine 14:*

The dialkenyl sulfonamide **13** (885 mg, 2.12 mmol, 1 eq.) and Grubbs catalyst (90 mg, 0.106 mmol, 5 mol%) in
10 degassed DCM (100 cm³) were heated to reflux for 18 h. The Grubb's catalyst used was tricyclohexylphosphine-[1,3-bis(2,4,6-trimethylphenyl)-4,5-dihydroimidazol-2-ylidene][benzylidene]ruthenium(IV)dichloride, available from Strum Chemicals Inc., Catalogue No. 77-7770. The
15 reaction mixture was cooled to room temperature and silica (ca. 3 g) was added. Solvent removal under reduced pressure and purification by flash column chromatography (Pet:EtOAc; 3:1) gave the title compound **14** (780 mg, 96%) as a clear viscous oil. $R_f = 0.25$
20 (Pet:EtOAc; 3:1); m/z (CI) 385 (MH^+ , 5%), 355 (20%), 198 (90%), 94 (100%); found 385.1224, $C_{20}H_{20}N_2O_4S \cdot H$ requires 385.1222 (+0.5 ppm); δ_H (300 MHz, $CDCl_3$) 1.72-1.82 (1H, m, CH_2), 1.83-2.04 (3H, m, CH_2), 2.13-2.24 (2H, m, CH_2), 3.49 (1H, pent, J 8.0 Hz, CH), 3.67 (2H, d, J 7.5 Hz, CH_2), 4.19 (2H, s, CH_2), 5.90 (1H, dt, J 2.0, 7.5 Hz, CH), 7.25-7.32 (2H, m, ArH), 7.35-7.42 (2H, m, ArH), 7.64-7.77 (3H, m, ArH), 8.05 (1H, dd, J 2.0, 5.5 Hz, ArH); δ_C (75 MHz, $CDCl_3$) 17.8, 28.4, 39.5,

43.0, 49.2, 116.9, 124.0, 126.1, 128.0, 129.8, 130.5,
131.5, 132.9, 133.2, 133.3, 139.9, 148.2, 151.0.

Example 15a



5

5-Cyclobutyl-2-(2-nitrophenylsulfonyl)-2,3,4,5-tetrahydro-1H-benzo[c]azepine-4,5-diol 15:

At room temperature the alkene 14 (120 mg, 0.313 mmol, 1 eq.) was dissolved in acetone (5 cm³) and H₂O (2.5 cm³) and NMO (40 mg, 0.341 mmol, 1.1 eq.) were added. OsO₄ (8 mg, 0.0315 mmol, 10 mol%) was then added to the vigorously stirred mixture. Stirring was continued for 18 h. DCM (15 cm³) and H₂O (15 cm³) were added and the mixture was acidified with 3 M HCl (pH 2). The aqueous phase was further extracted with DCM (3 x 15 cm³) and the combined organics were dried over MgSO₄. Filtration, solvent removal in vacuo followed by flash column chromatography (Pet:EtOAc; 2:1) gave 15 (104 mg, 80%) as an amorphous grey solid. The diol 15 was further purified by reprecipitation from Et₂O and petroleum ether. *R_f* = 0.3 (Pet:EtOAc; 1:1); *ν*_{max} (CDCl₃, cm⁻¹) 3540, 3093, 2982, 2940, 2867, 1590, 1545, 1445, 1371, 1352, 1164; *m/z* (CI) 436 (MNH₄⁺, 10%), 419 (MH⁺, 5%), 389 (20%), 232 (40%), 94 (100%); found 419.1274, C₂₀H₂₂N₂O₆S·H requires 419.1277 (-0.7 ppm); *δ*_H (300 MHz, CDCl₃) 1.29 (1H, m, CH₂), 1.78-1.90 (3H, m, CH₂), 2.13-2.38 (2H, m, CH₂), 2.49 (1H, d, *J* 9.25


Hz, OH), 2.91 (1H, pent, J 8.0 Hz, CH), 3.22 (1H, s, OH), 3.53 (1H, dd, J 1.0, 15.0 Hz, CH₂), 3.86 (1H, m, CH), 4.03 (1H, ddd, J 2.0, 4.0, 15.0 Hz, CH₂), 4.46 (1H, d, J 16.0 Hz, CH₂), 4.83 (1H, dd, J 2.0, 16.0 Hz, CH₂), 7.23-7.26 (2H, m, ArH), 7.36-7.40 (1H, m, ArH), 7.67-7.81 (3H, m, ArH), 7.86 (1H, d, J 7.5 Hz), 8.11 (1H, dd, J 2.0, 7.0 Hz, ArH); δ_C (75 MHz, CDCl₃) 17.5, 21.6, 21.8, 39.3, 50.9, 54.2, 72.5, 79.2, 124.2, 127.6, 128.2, 129.2, 130.2, 131.4, 131.7, 132.2, 132.8, 133.9, 140.8, 147.9.

Example 15b

5-Cyclobutyl-2-(2-nitrophenylsulfonyl)-2,3,4,5-tetrahydro-1H-benzo[*c*]azepine-4R,5R-diol 15:


At 5°C a mixture of dihydroazepine 14 (96 mg, 0.25 mmol, 1 eq.), AD-mix- β (180 mg) and MeSO₂NH₂ (24 mg, 0.25 mmol, 1 eq.) in *t*BuOH (1 cm³) and H₂O (1 cm³) were stirred for 2 days. Saturated Na₂SO₃ (5 cm³) and DCM (10 cm³) were added and the mixture was partitioned vigorously for 1 h. The resultant aqueous phase was further extracted with DCM (3 x 10 cm³) and the combined organic phases were dried over MgSO₄. Filtration, solvent removal and purification by flash column chromatography (Pet:EtOAc; 1:1) afforded 14 (41 mg, 43%) and the diol 15 (52 mg, 50%) whose data was in agreement to that reported above.

Example 16



5-Cyclobutyl-2,3,4,5-tetrahydrobenzo[c]azepine-4,5-diol
16:


At room temperature a mixture of nosylate 15 (256 mg, 0.61 mmol, 1 eq.) and K₂CO₃ (296 mg, 2.14 mmol, 3.5 eq.) in DMF (15 cm³) was treated with PhSH (80 µL, 0.78 mmol, 1.3 eq.). Stirring was continued at room temperature for 24 h. Ethyl acetate (25 cm³) and water (25 cm³) was added and the resultant aqueous layer was further extracted with EtOAc (5 x 25 cm³). The combined organic extracts were dried over MgSO₄. Filtration followed by solvent removal and column chromatography (EtOAc:MeOH; 3:1, 1% TEA) afforded the diol 16 (122 mg, 86%) as an amorphous colourless solid. R_F = 0.25 (EtOAc:MeOH; 3:1, 1% TEA); m/z (CI) 234 (MH⁺, 100%); found 234.1498, C₁₄H₁₉NO₂·H requires 234.1494 (+1.7 ppm); δ_H (300 MHz) 1.26-1.38 (1H, m, CH₂), 1.69-1.88 (3H, m, CH₂), 2.07-2.28 (2H, m, CH₂), 2.81-2.94 (1H, m, CH), 3.05 (1H, d, J 13.5 Hz, CH₂), 3.18 (1H, dd, J 3.5, 13.5 Hz, CH₂), 3.66 (1H, d, J 3.5 Hz, CH), 3.84 (1H, d, J 15.0 Hz, CH₂), 4.00 (1H, d, J 15.0 Hz, CH₂), 7.01 (1H, d, J 7.5 Hz, ArH), 7.16 (1H, t, J 7.5 Hz, ArH), 7.28 (1H, t, J 7.5 Hz, ArH), 7.80 (1H, d, J 7.5 Hz, ArH); δ_C (75 MHz) 17.6, 21.6, 21.8, 39.7, 51.9, 55.8, 73.2, 79.8, 126.9, 127.1, 129.2, 129.5, 136.3, 141.7.

Example 17

N-Benzyl-N-[2-(5-cyclobutyl-4,5-dihydroxy-1,3,4,5-tetrahydrobenzo[c]azepin-2-yl)ethyl]-2-nitrobenzenesulfonamide 17:

At room temperature **16** (122 mg, 0.524 mmol, 1 eq.) and the aldehyde **6** (see Example 6) (350 mg, 1.048 mmol, 2 eq.) in MeOH (5 cm) were treated with NaBH₃CN (33 mg, 0.525 mmol, 1 eq.) and conc. HCl (1 drop). The mixture was stirred for 15 h before EtOAc (25 cm³) and H₂O (25 cm³) were added. The pH was adjusted with 1 M NaOH to ca. 12 and the resultant aqueous layer was further extracted with EtOAc (2 x 25 cm³) and DCM (3 x 25 cm³). The combined organic extracts were dried over MgSO₄ and filtered. Silica (ca. 2.5 g) was added and the solvent was removed in vacuo. Purification by flash column chromatography (Pet:EtOAc; 1:1; 1% TEA → EtOAc; 1% TEA) gave the adduct **17** (189 mg, 66%) as a viscous yellow oil. *R_f* = 0.2 (Pet:EtOAc; 1:1); *ν*_{max} (CDCl₃/cm⁻¹) 3468, 3065, 2940, 2867, 1544, 1455, 1370, 1162; *m/z* (CI) 552 (MH⁺, 2%), 363 (10%), 108 (100%); found 552.2172, C₂₉H₃₃N₃O₆S·H requires 552.2168 (+0.7 ppm); δ _H (300 MHz, CDCl₃) 1.19-1.31 (1H, m, CH₂), 1.57-1.77 (3H, m, CH₂), 2.05-2.22 (2H, m, CH₂), 2.50-2.62 (2H, m, CH₂), 2.71 (1H, pent, *J* 8.5 Hz, CH), 2.82-2.93 (2H, m, CH₂), 3.31 (1H, ddd, *J* 5.0, 8.5, 15.0 Hz, CH₂), 3.38


(1H, dd, J 5.5, 15.0 Hz, CH₂), 3.44 (1H, dd, J 7.5, 15.0 Hz, CH₂), 3.58 (1H, d, J 3.5 Hz, CH), 3.69 (1H, d, J 15.0 Hz, CH₂), 6.91 (1H, d, J 7.5 Hz, ArH), 7.10 (1H, dt, J 1.5, 7.5 Hz, ArH), 7.18-7.31 (6H, m, ArH), 7.53-7.67 (3H, m, ArH), 7.73 (1H, dd, J 1.5, 8.0 Hz, ArH), 7.95 (1H, d, J 8.0 Hz, ArH); δ_C (100 MHz, CDCl₃) 17.7, 21.7, 21.8 (CH₂), 39.6 (CH), 45.5, 52.2, 57.5, 60.2, 63.0 (CH₂), 73.1 (CH), 79.3 (C-quat), 124.2, 126.8, 127.2, 128.1, 128.15, 128.7, 128.8, 130.2, 130.8, 131.6 (CH), 133.1 (C-ipso), 133.4 (CH), 134.5, 135.1, 141.4, 147.8 (C-ipso).

Example 18

15 N-Benzyl-N-[2-(5-cyclobutyl-5-hydroxy-4-oxo-1,3,4,5-tetrahydrobenzo[c]azepin-2-yl)ethyl]-2-nitrobenzenesulfonamide 18:

A solution of (COCl)₂ (42 μ L, 0.481 mmol, 4 eq.) in DCM (1 cm³) was treated at -78°C with DMSO (60 μ L, 0.846 mmol, 7 eq.). Stirring was continued for 0.25 h before a solution of 17 (60 mg, 0.109 mmol, 1 eq.) in DCM (1 cm³) was added in a dropwise fashion. Additionally, the flask was washed with DCM (1 cm³). The reaction mixture was stirred for 2 h during which time the temperature reached -10°C. TEA (100 μ L, 0.718 mmol, 6 eq.) was added to the reaction mixture. Stirring was

continued for x h before H₂O (20 cm³) and Et₂O (20 cm³) were added. The resultant aqueous layer was further extracted with Et₂O (4 x 20 cm³) and the combined organic extracts were dried over MgSO₄. Filtration followed by solvent removal and purification by flash column chromatography (Pet:EtOAc; 3:1, 1% TEA → Pet:EtOAc; 1:1, 1% TEA) afforded 18 (46 mg, 77%) as a yellow oil. $R_f = 0.2$ (Pet:EtOAc; 3:1, 1% TEA); ν_{\max} (neat/cm⁻¹) 3466, 3065, 2936, 2861, 1698, 1544, 1454, 1369, 1163; m/z (CI) 550 (MH⁺, 20%), 363 (40%), 106 (90%), 94 (100%); found 550.2021, C₂₉H₃₁N₃O₆S·H requires 550.2012 (+1.6 ppm); δ_H (300 MHz, CDCl₃) 1.49-1.62 (1H, m, CH₂), 1.65-1.88 (4H, m, CH₂), 2.22 (1H, pent, J 9.0 Hz, CH), 2.33 (2H, t, J 7.0 Hz, CH₂), 3.34-3.39 (3H, m, CH, CH₂), 3.38 (1H, d, J 15.0 Hz, CH₂), 3.64 (1H, d, J 15.0 Hz, CH₂), 3.76 (1H, d, J 16.0 Hz, CH₂), 4.04 (1H, d, J 16.0 Hz, CH₂), 4.50 (2H, s, CH₂), 4.54 (1H, s(br), OH), 6.91 (1H, d, J 7.5 Hz, ArH), 7.10 (1H, dt, J 1.5, 7.5 Hz, ArH), 7.18-7.31 (6H, m, ArH), 7.53-7.67 (3H, m, ArH), 7.73 (1H, dd, J 1.5, 8.0 Hz, ArH), 7.95 (1H, dd, J 1.0, 8.0 Hz, ArH); δ_C (75 MHz, CDCl₃) 17.0, 21.3, 21.6, 41.6, 44.8, 51.0, 52.1, 59.5, 63.5, 84.1, 124.2, 127.3, 127.5, 127.6, 128.1, 128.2, 128.7, 129.9, 130.8, 131.6, 133.4, 133.6, 133.8, 135.4, 138.3, 148.1, 206.8.


Example 19

2-[2-(benzylamino)ethyl]-5-cyclobutyl-5-hydroxy-
1,3,4,5-tetrahydrobenzo[c]azepin-4-one **19**:

- 5 At room temperature a solution **18** (260 mg, 0.474 mmol, 1 eq.) in DMF (5 cm³) was treated with K₂CO₃ (212 mg, 1.534 mmol, 3.2 eq.) and PhSH (60 µL, 0.584 mmol, 1.2 eq.). Stirring was continued for 18 h before H₂O (25 cm³) and EtOAc (25 cm³) were added. The resultant
- 10 aqueous layer was extracted with EtOAc (5 x 15 cm³) and the combined organic layers were dried over MgSO₄. The crude amine obtained after filtration and solvent removal *in vacuo* was purified by flash column chromatography (Pet:EtOAc; 1:1, 1% TEA → Pet:EtOAc;
- 15 1:2, 1% TEA) which gave the title compound **19** (90 mg, 52%) as a yellow oil. *R*_f = 0.15 (Pet:EtOAc; 1:1, 1% TEA); *ν*_{max} (neat/cm⁻¹) 3454, 3054, 2934, 2855, 1692, 1453; *m/z* (CI) 365 (MH⁺, 80%), 347 (M-OH⁺, 30%), 108 (100%), 74 (80%); found 365.2220, C₂₃H₂₈N₂O₂·H requires
- 20 365.2229 (-2.5 ppm); *δ*_H (300 MHz, CDCl₃) 1.40-1.48 (1H, m, CH₂), 1.58-1.76 (4H, m, CH₂), 2.08-2.16 (1H, m, CH₂), 2.48 (2H, t, *J* 6.0 Hz, CH₂), 2.71 (2H, t, *J* 6.0 Hz, CH₂), 3.26 (1H, pent, *J* 8.5 Hz, CH), 3.44 (1H, d, *J* 15.5 Hz, CH₂), 3.75 (2H, d, *J* 15.5 Hz, CH₂), 3.76 (1H, d, *J* 16.5 Hz, CH₂), 3.90 (1H, d, *J* 13.5 Hz, CH₂), 3.96
- 25 (1H, d, *J* 13.5 Hz, CH₂), 4.15 (1H, d, *J* 16.5 Hz, CH₂),

6.84 (1H, d, J 7.5 Hz, ArH), 7.05 (1H, dt, J 1.0, 7.5 Hz, ArH), 7.11-7.36 (6H, m, ArH), 7.64 (1H, d, J 7.5 Hz, ArH).

5 Example 20




N-Benzyl-N-[3-(5-cyclobutyl-4,5-dihydroxy-1,3,4,5-tetrahydrobenzo[c]azepin-2-yl)propyl]-2,2,2-trifluoroacetamide **20**:

- 10 At room temperature the nosylate **15** (120 mg, 0.287 mmol, 1 eq.) and K_2CO_3 (129 mg, 0.933 mmol, 3.25 eq.) in DMF (5 cm³) was treated with phenylmercaptan (44 μ L, 0.428 mmol, 1.49 eq.). Stirring was continued for 24 h before the reaction mixture was exhaustively extracted
- 15 with EtOAc (5 x 25 cm³) and H₂O (25 cm³). The combined organic extracts were dried over $MgSO_4$, filtration followed by solvent removal *in vacuo* afforded the crude diol **16**. A mixture of the crude diol **16** (ca. 0.287 mmol, 1 eq.) the aldehyde **3** (223 mg, 0.861 mmol, 3
- 20 eq.), $NaBH_3CN$ (18 mg, 0.286 mmol, 0.99 eq.) in MeOH (5 cm³) with a drop of conc. HCl were stirred at room temperature for 15 h. The reaction mixture was extracted with Et₂O (5 x 25 cm³) and 1 M NaOH (25 cm³), dried over $MgSO_4$. Filtration followed by solvent
- 25 removal under reduced pressure afforded the crude adduct which was purified by column chromatography (Pet:EtOAc; 3:1, 1% TEA \rightarrow Pet:EtOAc; 1:1, 1% TEA) gave the title compound **20** (104 mg, 76%). R_f = 0.1 (Pet:EtOAc; 3:1, 1% TEA); 0.3 (Pet:EtOAc; 1:1, 1%

TEA); m/z (CI) 477 (MH^+ , 100%); found (EI) 476.2287, $C_{26}H_{31}N_2O_3F_3$ requires 476.2290 (-0.8 ppm); δ_H (300 MHz, $CDCl_3$) 1.28-1.45 (1H, m, CH_2), 1.63-1.91 (3H, m, CH_2), 2.14-2.36 (2H, m, CH_2), 2.45-2.59 (2H, m, CH_2), 2.61-2.72 (2H, m, CH_2), 2.75-2.96 (2H, m, CH_AH_B , CH), 2.98-3.05 (1H, m, CH_AH_B), 3.22-3.42 (2H, m, CH_2), 3.50-3.61 (1H, m, CH_AH_B), 3.62-3.75 (1H, m, CH), 3.75-3.82 (1H, m, CH_AH_B), 4.52 $^{\wedge}$ (1H, d, J 14.5 Hz, CH_2), 4.56* (1H, d, J 15.5 Hz, CH_2), 4.69* (1H, d, J 15.5 Hz, CH_2), 4.82 $^{\wedge}$ (1H, d, J 14.5 Hz, CH_2), 7.05-7.18 (2H, m, ArH), 7.19-7.25 (1H, m, ArH), 7.25-7.45 (5H, m, ArH), 7.80-7.88 (1H, m, ArH). 1H -NMR complicated due to rotameric structures [60*:40 $^{\wedge}$].

15 Example 21




N-Benzyl-N-[3-(5-cyclobutyl-5-hydroxy-4-oxo-1,3,4,5-tetrahydrobenzo[c]azepin-2-yl)propyl]-2,2,2-trifluoroacetamide **21**:

20 A solution of $(COCl)_2$ (98 μL , 1.12 mmol, 3 eq.) in DCM (3 cm^3) was treated at $-78^\circ C$ with DMSO (133 μL , 1.87 mmol, 5 eq.). Stirring was continued for 10 min. before a solution of **20** (179 mg, 0.376 mmol, 1 eq.) in DCM (3 cm^3) was added in a dropwise fashion. This
25 flask was washed with DCM (2 cm^3) and this was also added. The reaction mixture was stirred for 1 h during which time the temperature reached $0^\circ C$. TEA (0.31 cm^3 ,

2.22 mmol, 6 eq.) was added to the reaction mixture and stirring was continued for 0.5 h. H₂O (25 cm³) and Et₂O (25 cm³) were added and the resultant aqueous layer was further extracted with Et₂O (4 x 25 cm³) and the combined organic extracts were dried over MgSO₄. Filtration followed by solvent removal and purification by flash column chromatography (Pet:EtOAc; 5:1, 1% TEA) afforded **21** (120 mg, 67%) as a yellow oil. *R*_f = 0.3 (Pet:EtOAc; 5:1, 1% TEA); *ν*_{max} (neat/cm⁻¹) 3466, 3065, 2942, 2861, 1691, 1452, 1376, 1202, 1144; *m/z* (CI) 475 (MH⁺, 5%), 261 (40%), 221 (80%), 108 (100%), 91 (90%); found 475.2205, C₂₆H₂₉N₂O₃F₃·H requires 475.2208 (-0.6 ppm); ¹H (200 MHz, CDCl₃) 1.49-1.98 (8H, m, CH₂), 2.18-2.40 (2H, m, CH₂), 3.28-3.60 (4H, m, CH₂, CH_AH_B, CH), 3.74-3.90 (2H, m, CH_AH_B, CH_A'H_B'), 4.15-4.31 (1H, m, CH_A'H_B'), 4.55-4.73 (2H, m, CH₂), 6.95-7.01 (1H, m, ArH), 7.10-7.45 (7H, m, ArH), 7.71-7.83 (1H, m, ArH). ¹H-NMR complicated due to rotameric structures.

20 Example 22



2-[3-(benzylamino)propyl]-5-cyclobutyl-5-hydroxy-1,2,3,5-tetrahydrobenzo[c]azepin-4-one **22**:

At room temperature a solution of **21** (80 mg, 0.169 mmol, 1 eq.) in MeOH (10 cm³) was treated with a solution of K₂CO₃ (117 mg, 0.847 mmol, 5 eq.) in H₂O (0.6 cm³). Stirring was maintained for 24 h. Solvent removal in vacuo followed by purification by flash

column chromatography (Pet:EtOAc; 1:2, 1% TEA) gave the title compound 22 (50 mg, 80%) as a yellow oil. R_f = 0.2 (Pet:EtOAc; 1:2); m/z (CI) 379 (MH^+ , 50%), 284 (50%), 267 (55%), 108 (100%); found 379.2381, $C_{24}H_{30}N_2O_2 \cdot H$ requires 379.2385 (-1.1 ppm); δ_H (300 MHz, $CDCl_3$) 1.41-1.88 (7H, m, CH_2), 2.13-2.28 (1H, m, CH_2), 2.38 (2H, dt, J 3.5, 7.0 Hz, CH_2), 2.59 (2H, t, J 7.0 Hz, CH_2), 3.39-3.51 (1H, m, CH), 3.45 (1H, d, J 15.75 Hz, CH_2), 3.68 (1H, d, J 15.75 Hz, CH_2), 3.69 (2H, s, CH_2), 3.85 (1H, d, J 16.25 Hz, CH_2), 4.14 (1H, d, J 16.25 Hz, CH_2), 6.97 (1H, d, J 7.5 Hz, ArH), 7.11 (1H, dt, J 1.5, 7.5 Hz, ArH), 7.14-7.26 (6H, m, ArH), 7.67 (1H, d, J 7.5 Hz, ArH); δ_C (75 MHz, $CDCl_3$) 17.2, 21.6, 21.7, 27.4, 30.3, 41.7, 47.1, 52.5, 60.1, 64.1, 84.0, 127.0, 127.3, 127.4, 128.0, 128.4, 129.6, 134.5, 138.4, 207.7.

Pharmacology

20 Functional assays of M1, M2 and M3 receptor activity

Initial evaluation of test compounds is by assay of functional tissue responses. This has the advantage that it readily discriminates between agonist partial agonist and antagonist activity

25 M1 - Vas deferens preparations

Male New Zealand white rabbits (1.47 - 3.4 kg) are killed by a blow to the back of the head and vasa deferentia removed, dissected free of connective tissue and divided into prostatic and epididymal portions.

30 Each segment is mounted on a tissue holder and passed through two ring electrodes (5mm apart). They are immersed in a modified low Ca^{2+} Krebs solution at

32±0.51°C and gassed with 5% CO₂ in oxygen. Yohimbine (1.0mM) is present throughout to block prejunctional α₂-adrenoceptors. The upper end of the tissue is attached by cotton thread to an isometric transducer (MLT020, ADInstruments). Tissues are left to equilibrate for at least 45 min at passive force of 0.75-1g. Field stimulation is then applied by repeated application of single pulses (30V, 0.05Hz, 0.5ms). Isometric tension is recorded by computer at a sampling rate of 100Hz, using Powerlab/200 (ADInstruments) software and MacLab bridge amplifiers.

M2 - Guinea-pig atria

Guinea-pigs are killed by a blow to the back of the head and left atrium removed. The atrium is secured to a pair of stainless steel electrodes by means of a cotton thread and immersed in the organ bath containing gassed Krebs solution with normal Ca²⁺ at 32±0.5°C. Atria are placed at 2Hz with square-wave pulses of 0.5ms pulse width. Isometric contractions are recorded by computer or polygraph.

M3 - Guinea-pig ileum

Sections (2 cm) are cut from the ileum of the killed guinea-pigs, 10cm from the ileo-caecal junction. One end is attached to a tissue holder/aerator and the other end via a cotton thread to an isometric transducer. The tissue is immersed in gassed normal Ca²⁺ Krebs solution at 32±0.5°C. A resting tension of 0.5g is applied and isometric contractions measured by computer or polygraph.

30 Agonist concentration-response curves

Following at least 30 min equilibration to allow twitches or tension to stabilize, cumulative concentration-response curves for the muscarinic

agonists are constructed. The concentration is increased in half logarithmic increments after the contraction in the presence of each concentration has plateaued. Steady-state contractions at each concentration are measured and the inhibition expressed as a percentage of the baseline twitch height in atria and vas deferens or as the maxi contraction in the ileum. EC50 values for the muscarinic agonists are determined from individual curves as the molar concentration required for 50% inhibition of twitch height or the 50% of maximum contraction (ileum). Geometric mean EC50 values and their 95% confidence limits are calculated.

Effects of muscarinic antagonists

A concentration-response curve to the test agonist is established in the absence of antagonist and after achieving the maximum effect, the agonist washed from the bath to restore twitch contractions. Three further concentration-response curves are then obtained in the same manner at approximately 30 min intervals, with the antagonist (Standards - pirenzepine M1, darifenacin M3, methoctramine M2) being introduced to the bath 15 min before each of these subsequent curves.

Calculation of antagonist pA_2 values

Concentration-response curves in the absence and presence of antagonist are measured as described for the agonist studies. The shifts in the concentration-response curves in the presence of antagonist compared with the absence of antagonist are expressed as the dose-ratios (DR) of the EC50 values. pA_2 values are then determined from Schild analysis of plots of the mean corrected $-\log(DR-1)$ against log molar concentration of antagonist. The slopes of the Schild

plots are determined by linear regression and the pA_2 values determined from the intercept on the concentration axis (when $\log(DR-1)$ is zero). pA_2 values are also determined from individual concentrations of antagonist by applying the equations: $pA_2 = \log(DR-1) = \log[B]$, wherein B is the molar concentration of antagonist.

Standard Drugs

Carbamoylcholine chloride (carbachol),
10 methacholine, methacramine, pirenzepine
dihydrochloride, yohimbine hydrochloride (Sigma, Poole, Dorset, UK), darifenacin (Pfizer, Sandwich, Kent), McN-A343 [4-(4-chlorophenylcarbamoyloxy)-2-butyryltrimethylammonium iodide] and oxotremorine
15 sesquifurnarate (RBI, St. Albans, UK). AR drugs are dissolved in distilled water initially and dilutions made in Krebs solution.

Reference data


The reproducibility of the concentration-response
20 curves and stability of each tissue over several hours was established. EC50 values were obtained for a range of reference agonists in each tissue - methacholine, oxotremorine, McN-A-343 to permit comparisons with the novel agents of the invention. pA_2 values for reference
25 antagonists were obtained in relevant tissues - pirenzepine (M1 selective), darifenacin (M3 selective). It is as a result possible to establish the functional characterization of the three receptor types to enable determination of the potency (EC50 of agonist, pA_2 or
30 affinity of antagonist molecules) and selectively of the novel agents of the invention.

The compounds of Examples 19 and 22 were tested as described above and the results obtained were as follows:

Compound	M3 (ileum)	M3 (Atria)	Log ₁₀ selectivity
Compound 22 (Example 22)	6.7±0.4 (4 pts)	5.2±0.3 (4 pts)	1.5 ± 0.5
Compound 19 (Example 19)	6.5±0.4 (3 pts)	4.9±0.6 (4 pts)	1.6 ± 0.7

CLAIMS:

1. A compound having the formula:




wherein:


R_{1a} , R_{1b} and R_{1c} are independently fluorine or hydrogen;

R_2 is C_1 to C_{12} alkyl, said alkyl being straight or branched chain, saturated or unsaturated, mono-substituted or unsubstituted, said substituents being selected from piperidine, pyrrolidine, morpholine, thiomorpholine, tetrahydrofuran, thiophen, furan and cycloalkyl of 3 to 7 carbon atoms; a cycloalkyl of 3 to 9 carbon atoms; a cycloalkyl of 3 to 9 carbon atoms (preferably 4 to 9 carbon atoms) having a C_1 to C_6 alkyl substituent; a polycycloalkyl of 2 to 3 rings having 7 to 12 carbons; and phenyl or phenyl singly or multiply substituted (preferably singly or doubly) with halogen, hydroxy, C_1 to C_6 alkoxy, C_1 to C_6 alkyl, nitro, methylene dioxy or trifluoromethyl; and

R_3 is a moiety selected from:




I



II

or a pyrrolidin-3-yl moiety of the formula

-55-



III

where R₆ is hydroxy or hydrogen;

where one of R₄ and R₅ is hydrogen or lower C1-3 alkyl
and the other is selected from:

- (a) hydrogen,
 - (b) phenyl,
 - (c) phenyl singly or multiply substituted with halogen, hydroxy, C₁ to C₆ alkoxy, C₁ to C₆ alkyl, nitro, methylene dioxy or trifluoromethyl,
 - (d) C₁ to C₆ alkyl which may be branched chain or straight, saturated, unsaturated, or cyclic and may be optionally substituted with hydroxy, thienyl, pyrrolyl, pyridyl, furanyl, lower alkoxy or acetoxyalkyl wherein the alkyl group has 1 to 3 carbons, phenyl, phenyl singly or multiply substituted (preferably singly or doubly) with halogen, hydroxy, C₁ to C₁₀ alkoxy, C₁ to C₁₀ alkyl, nitro, methylene dioxy (optionally mono or di-alkyl substituted where the alkyl substituent has from 1 to 10 carbon atoms) or trifluoromethyl;
- or a pharmaceutically acceptable salt thereof
2. A compound according to claim 1, wherein R₂ is cycloalkyl of 3 to 6 carbon atoms.
 3. A compound according to claim 2, wherein R₂ is cyclobutyl.
 4. A compound according to any preceding claim, wherein R₄ is hydrogen and R₅ is selected from amongst the groups (a)-(d) as defined in claim 1.
 5. A compound according to any one of claims 1 to 3, wherein one of R₄ and R₅ is hydrogen (or methyl in the case of R₅) and the other is selected from hydrogen,

-56-

C₁ to C₆ alkyl which may be branched chain or straight, saturated, unsaturated, or cyclic and may be optionally substituted with hydroxy, thienyl, pyrrolyl, pyridyl, furanyl, phenyl, phenyl singly or multiply substituted (preferably singly or doubly) with halogen, hydroxy, C₁ to C₁₀ alkoxy, C₁ to C₁₀ alkyl or nitro.


6. A compound according to claim 5, wherein R₄ is hydrogen and R₅ is C₁ to C₆ alkyl substituted by phenyl or phenyl which is singly or multiply substituted with halogen, hydroxy, C₁ to C₁₀ alkoxy, C₁ to C₁₀ alkyl or nitro.

7. A compound according to claim 6, wherein R₅ is benzyl, substituted benzyl or cinnamyl.

8. A compound according to claim 7, wherein R₅ is substituted benzyl in which the substituent(s) on the benzyl are independently halo, C₁ to C₁₀ alkoxy or C₁ to C₁₀ alkyl.

9. A compound according to any preceding claim, wherein R₆ is hydrogen.

10. A compound according to claim 1, wherein R_{1a}, R_{1b} and R_{1c} are independently hydrogen or fluorine, R₂ is cycloalkyl of 3 to 6 carbon atoms or phenyl, R₃ is




where R₄ is hydrogen and R₅ is selected from C₁ to C₆ alkyl, benzyl, substituted benzyl or cinnamyl, and R₆ is hydrogen or hydroxy.

11. A compound according to claim 10, wherein R₂ is cyclobutyl and R₅ is substituted benzyl in which the substituent(s) on the benzyl are independently halo, C₁ to C₁₀ alkoxy or C₁ to C₁₀ alkyl.

-57-

12. A compound according to claim 10 or 11,
wherein R_6 is hydrogen.

13. A compound according to claim 1, wherein R_{1a} ,
 R_{1b} and R_{1c} are independently hydrogen or fluorine, R_2 is
5 cycloalkyl of 3 to 6 carbon atoms or phenyl, R_3 is




10 where R_4 is hydrogen and R_5 is selected from C_1 to C_6
alkyl, benzyl, substituted benzyl or cinnamyl, and R_6 is
hydroxy or hydrogen.

14. A compound according to claim 13, wherein R_2
15 is cyclobutyl and R_5 is substituted benzyl in which the
substituent(s) on the benzyl are independently halo, C_1
to C_{10} alkoxy or C_1 to C_{10} alkyl.

15. A compound according to claim 13 or 14,
wherein R_6 is hydrogen.

16. A compound according to claim 15, wherein R_{1a} ,
20 R_{1b} and R_{1c} are independently hydrogen or fluorine, R_2 is
cycloalkyl of 3 to 6 carbon atoms or phenyl, R_3 is a
moiety having the following structure:



25 where R_4 is hydrogen and R_5 is selected from C_1 to C_6
alkyl, benzyl, substituted benzyl or cinnamyl.

17. A compound according to claim 16, wherein R_2
is cyclobutyl and R_5 is substituted benzyl in which the
substituent(s) on the benzyl are independently halo, C_1
35 to C_{10} alkoxy or C_1 to C_{10} alkyl.

-58-

18. A compound according to any preceding claim, wherein R_{1a} , R_{1b} and R_{1c} are each hydrogen.

19. A compound according to claim 1, which is 2-[2-(benzylamino)ethyl]-5-cyclobutyl-5-hydroxy-1,3,4,5-tetrahydrobenzo[c]azepin-4-one, or a pharmaceutically acceptable salt thereof.


20. A compound according to claim 1, which is 2-[3-(Benzylamino)propyl]-5-cyclobutyl-5-hydroxy-1,2,3,5-tetrahydrobenzo[c]azepin-4-one, or a pharmaceutically acceptable salt thereof.

21. A pharmaceutical composition comprising a compound according to any preceding claim and a pharmaceutically acceptable carrier or diluent.

22. A compound as claimed in any one of claims 1 to 20, for use as a muscarinic antagonist with M_3 selectivity.

23. A compound for use as claimed in claim 22, as a bronchodilator, an antispasmodic agent, an antisecretory agent, an agent having antiulcer activity or a agent for the treatment of patients suffering from neurogenic bladder disorders.

24. A process for synthesising a compound according to claim 1, which includes the step of subjecting a compound of the formula (X):



in which R_{1a} , R_{1b} , R_{1c} and R_2 are as defined in claim 1 and R_3 is as defined in claim 1 suitably protected to oxidation conditions sufficient to oxidise the alcohol group at the 4-position of the benzo[c]azepine core to a ketone group.

INTERNATIONAL SEARCH REPORT

Interr	ication No
PCT/GB 01/02594	

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 C07D223/16 A61K31/55 A61P11/08 A61P13/10 A61P1/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

PAJ, EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PATENT ABSTRACTS OF JAPAN vol. 1996, no. 02, 29 February 1996 (1996-02-29) & JP 07 258250 A (YAMANOUCHI PHARMACEUT CO LTD), 9 October 1995 (1995-10-09) abstract -----	1-24

☐ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *S* document member of the same patent family

Date of the actual completion of the international search

30 August 2001

Date of mailing of the international search report

10/09/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Schmid, J-C

INTERNATIONAL SEARCH REPORT

Intern	ication No
PCT/GB 01/02594	

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
JP 07258250 A	09-10-1995	NONE	
